Author Index

Adachi, K., see Yamaga, N. 25
Amagaya, S., see Umeda, M. 246
Araki, A.
– and Sako, Y.
Determination of free and total homocysteine in human plasma by high-performance liquid chromatography with fluorescence detection 43
Bandô, Y., see Yamaguchi, Y. 53
Bayne, W.F., see Chiou, R. 281
Beightol, L.A., see Karnes, H.T. 165
Biseth, A., see Christophersen, A.S. 117
Blackman, G.L., see Sitaram, B.R. 13
Blumer, J.L., see Myers, C.M. 153
Brault, D., see Dellinger, M. 73
Brons, P.P.T.
–, Wessels, J.M.C., Linssen, P.C.M., Haanen, C. and Speth, P.A.J.
Determination of amsacrine in human nucleated hematopoietic cells 175
Chang, D., see Oldfield, N. 135
Chiou, R.
–, Stubbs, R.J. and Bayne, W.F.
Improved method for the determination of theophylline in plasma and urine using high-performance liquid chromatography 281
Christophersen, A.S.
–, Biseth, A., Skuterud, B. and Gadeholt, G.
Identification of opiates in urine by capillary column gas chromatography of two different derivatives 117
Claycamp, H.G.
– and Ludwig, B.
Simplified monitoring of reduced and oxidized dithiothreitol using high-performance liquid chromatography 239
Crevat, A., see Gallice, P. 263
Davis, A.T.
Assay for trimethyllysine hydroxylase by high-performance liquid chromatography 253
Decolin, D., see Motassim, N. 340
Dellinger, M.
– and Brault, D.
Normal-phase high-performance liquid chromatography of free acid dicarboxylic porphyrins and hematoporphyrin derivative on silica 73
Denouel, J., see Humbert, H. 205
Desiderio, D.M., see Liu, D. 61
Dietel, P.
–, Hölzel, C., Schneider, K. and Spiteller, G.
Excretion of urinary acids during inverted sleep–waking rhythm 1
Dinh, T. Le, see Motassim, M. 340
Donaldson, J., see Workman, P. 315
Falkowski, A.J.
–, Look, Z.M., Noguchi, H. and Silber, B.M.
Determination of cefixime in biological samples by reversed-phase high-performance liquid chromatography 145
Farthing, D., see Karnes, H.T. 165
Gadeholt, G., see Christophersen, A.S. 117
Gall, J.-M. Le, see Gallice, P. 263
Gallice, P.
–, Monti, J.-P., Le Gall, J.-M., Crevat, A. and Murisasco, A.
Ion-pair high-performance liquid chromatography profiling of a uremic toxin fraction 263
Garland, W., see Oldfield, N. 135
Greenberg, J., see Phillips, M. 235
Greenblatt, D.J., see Kennedy, J.S. 274
Guyonnaud, C., see Lacroix, C. 217
Haanen, C., see Brons, P.P.T. 175
Haginaka, J., see Yamaguchi, Y. 53
Harmatz, J.S., see Kennedy, J.S. 274
Hayasaka, Y., see Sitaram, B.R. 13
Hölzel, C., see Dietel, P. 1
Hoffman, N.E., see Rustum, A.M. 125
Humbert, H.
–, Denouel, J. and Keller, H.P.
Column liquid chromatographic determination of hydrolysed bopindolol, in the picogram per millilitre range in plasma, using cartridge extraction and dual electrochemical detection 205
Inoue, T.
– and Suzuki, S.-I.
High-performance liquid chromatographic determination of triazolam and its metabolites in human urine 197
Iwai, K., see Shibata, K. 257
Karnes, H.T.
–, Opong-Mensah, K., Farthing, D. and Beightol, L.A.
Automated solid-phase extraction and high-performance liquid chromatographic determination of ranitidine from urine, plasma and peritoneal dialysate 165

Kawada, T., see Shibata, K. 257
Keller, H.P., see Humbert, H. 205
Kennedy, J.S.
–, Leduc, B.W., Scavone, J.M., Harmatz, J.S., Shader, R.I. and Greenblatt, D.J.
Pharmacokinetics of intravenous caffeine: comparison of high-performance liquid chromatographic and gas chromatographic methods 274
Kleinjans, J.C.S., see Verhagen, H. 288
Klockowski, P.M.
– and Levy, G.
Simultaneous determination of diazepam and its active metabolites in rat serum, brain and cerebrospinal fluid by high-performance liquid chromatography 334
Kohara, H., see Yamaga, N. 25
Koshakji, R.P.
– and Wood, A.J.J.
Improved high-performance liquid chromatographic method for the simultaneous determination of propranolol and 4-hydroxypropranolol in plasma with fluorescence detection 294
Kunimoto, M., see Yamaguchi, Y. 53
Lacroix, C.
–, Poncet, P., Laine, G., Guyonnaud, C., Ray, M., Menager, S. and Lafont, O.
Microdétermination du pyrazinamide et de ses métabolites (acide pyrazinoïque, acide hydroxy-5-pyrazinoïque, hydroxy-5-pyrazinamide, acide pyrazinurique) plasmatiques et urinaires par chromatographie en phase liquide 217
Lafont, O., see Lacroix, C. 217
Laine, G., see Lacroix, C. 217
Le Dinh, T., see Motassim, N. 340
Leduc, B.W., see Kennedy, J.S. 274
Lee, F.Y.F., see Workman, P. 315
Le Gall, J.-M., see Gallice, P. 263
Levy, G., see Klockowski, P.M. 334
Linssen, P.C.M., see Brons, P.P.T. 175
Liu, D.
– and Desiderio, D.M.
Reversed-phase high-performance liquid chromatographic–radioreceptor assay of human cerebrospinal fluid neuropeptides. Maximizing recovery of picomoles of peptides and minimizing memory 61
Lockett, L., see Sitaram, B.R. 13
Look, Z.M., see Falkowski, A.J. 145
Lora-Tamayo, C.
–, Tena, T. and Tena, G.
Concentrations of free and conjugated morphine in blood in twenty cases of heroin-related deaths 267
Ludwig, B., see Claycamp, H.G. 239
McLeish, M., see Sitaram, B.R. 13
McLeod, W.R., see Sitaram, B.R. 13
Maniwa, M., see Marunaka, T. 227
Marunaka, T.
– and Maniwa, M.
High-performance liquid chromatographic determination of [3,4-di-(4-methoxyphenyl)-5-isoxazolyl]acetic acid and its metabolites in human plasma and urine 227
Matsunaga, H.
–, Suehiro, T., Saita, T., Nakano, Y., Mori, M., Takata, K. and Oda, K.
High-performance liquid chromatographic determination of pyridostigmine in plasma 353
Maurer, H.
– and Pfleger, K.
Identification and differentiation of benzodiazepines and their metabolites in urine by computerized gas chromatography–mass spectrometry 85
Meffin, P.J., see Sallustio, B.C. 33
Menager, S., see Lacroix, C. 217
Mori, J.-P., see Gallice, P. 263
Mori, M., see Matsunaga, H. 353
Motassim, N.
–, Decolin, D., Le Dinh, T., Nicolas, A. and Siest, G.
Direct determination of dextromethorphan and its three metabolites in urine by high-performance liquid chromatography using a precolumn switching system for sample clean-up 340
Murisasco, A., see Gallice, P. 263
Myers, C.M.
– and Blumer, J.L.
High-performance liquid chromatography of ciprofloxacin and its metabolites in serum, urine and sputum 153
Nakano, Y., see Matsunaga, H. 353
Nicolas, A., see Motassim, M. 340
Noguchi, H., see Falkowski, A.J. 145
Norén, K.
– and Sjövall, J.
Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex® 5000 103
Oda, K., see Matsunaga, H. 353

Ogihara, Y., see Umeda, M. 246
Oldfield, N.
–, Chang, D., Garland, W. and Town, C.
Quantitation of ceftetrame in human plasma and urine by high-performance liquid chromatography 135
Opong-Mensah, K., see Karnes, H.T. 165
Oppitz, M., see Workman, P. 315
Pfleger, K., see Maurer, H. 85
Phillips, M.
– and Greenberg, J.
Detection of endogenous acetone in normal human breath 235
Pieters, F.A.J.M.
–, Vincken, B.J. and Zuidema, J.
Dapsone and monoacetyldapsone determined in serum and saliva by a sensitive high-performance liquid chromatographic method with a single extraction step 322
Poncet, P., see Lacroix, C. 217
Ray, M., see Lacroix, C. 217
Reiff, K.
High-performance liquid chromatographic method for the determination of adibendan in plasma and urine 346
Rustum, A.M.
– and Hoffman, N.E.
Determination of cyclophosphamide in whole blood and plasma by reversed-phase high-performance liquid chromatography 125
Saita, T., see Matsunaga, H. 353
Sako, Y., see Araki, A. 43
Sallustio, B.C.
–, Meffin, P.J. and Thompson, M.
High-performance liquid chromatographic quantitation of triacylglycerols containing fenoprofen from biological samples 33
Scavone, J.M., see Kennedy, J.S. 274
Schneider, K., see Dietel, P. 1
Shader, R.I., see Kennedy, J.S. 274
Shibata, K.
–, Kawada, T. and Iwai, K.
High-performance liquid chromatographic determination of nicotinamide in rat tissue samples and blood after extraction with diethyl ether 257
Shimizu, K., see Yamaga, N. 25
Siest, G., see Motassim, M. 340
Silber, B.M., see Falkowski, A.J. 145
Sitaram, B.R.
–, Lockett, L., McLeish, M., Hayasaka, Y., Blackman, G.L. and McLeod, W.R.
Gas chromatographic–mass spectroscopic characterisation of the psychotomimetic indolealkylamines and their in vivo metabolites 13
Sjövall, J., see Norén, K. 103
Skuterud, B., see Christophersen, A.S. 117
Soine, P.J.
– and Soine, W.H.
High-performance liquid chromatographic determination of the diastereomers of 1-(β-D-glucopyranosyl)amobarbital in urine 309
Soine, W.H., see Soine, P.J. 309
Soldin, S.J., see Tan, Y.K. 187
Sonneveld, P., see Van der Steuijt, K. 328
Speth, P.A.J., see Brons, P.P.T. 175
Spino, M., see Strong, H.A. 301
Spiteller, G., see Dietel, P. 1
Steuijt, K. van der, see Van der Steuijt, K. 328
Strong, H.A.
– and Spino, M.
Highly sensitive determination of cimetidine and its metabolites in serum and urine by high-performance liquid chromatography 301
Stubbs, R.J., see Chiou, R. 281
Suehiro, T., see Matsunaga, H. 353
Suzuki, S.-I., see Inoue, T. 197
Takata, K., see Matsunaga, H. 353
Tan, Y.K.
– and Soldin, S.J.
Analysis of salbutamol enantiomers in human urine by chiral high-performance liquid chromatography and preliminary studies related to the stereoselective disposition kinetics in man 187
Tena, G., see Lora-Tamayo, C. 267
Tena, T., see Lora-Tamayo, C. 267
Thijssen, H.H.W., see Verhagen, H. 288
Thompson, M., see Sallustio, B.C. 33
Town, C., see Oldfield, N. 135
Umeda, M.
–, Amagaya, S. and Ogihara, Y.
Separation of thromboxane B_2 by high-performance liquid chromatography 246
Van der Steuijt, K.
– and Sonneveld, P.
Concurrent analysis of methotrexate, trimethophrim, sulphamethoxazole and their major metabolites in plasma by high-performance liquid chromatography 328
Verhagen, H.
–, Thijssen, H.H.W. and Kleinjans, J.C.S.
Determination of butylated hydroxytoluene in plasma by high-performance liquid chro-

matography 288
Vincken, B.J., see Pieters, F.A.J.M. 322
Wessels, J.M.C., see Brons, P.P.T. 175
Wood, A.J.J., see Koshakji, R.P. 294
Workman, P.
–, Oppitz, M., Donaldson, J. and Lee, F.Y.F. High-performance liquid chromatography of chlorambucil analogues 315
Yamaga, N.
–, Adachi, K., Kohara, H. and Shimizu, K. Internal standards for quantitative gas chromatography of individual bile acids after group separation of bile acids in urine 25
Yamaguchi, Y.
–, Haginaka, J., Kunimoto, M., Yasuda, H. and Bandô, Y. High-performance liquid chromatographic determination of desmosine and isodesmosine in tissues and its application to studies of alteration of elastin induced by atherosclerosis 53
Yasuda, H., see Yamaguchi, Y. 53
Zuidema, J., see Pieters, F.A.J.M. 322

Subject Index

Acetone
Detection of endogenous acetone in normal human breath 235
N-Acetylsulphametoxazole
Concurrent analysis of methotrexate, trimethoprim, sulphamethoxazole and their major metabolites in plasma by HPLC 328
Adibendan
HPLC method for the determination of adibendan in plasma and urine 346
Amino acids
Excretion of urinary acids during inverted sleep–waking rhythm 1
Amobarbital
HPLC determination of the diastereomers of 1-(β-D-glucopyranosyl)amobarbital in urine 309
Amsacrine
Determination of amsacrine in human nucleated hematopoietic cells 175
Benzodiazepines
Identification and differentiation of benzodiazepines and their metabolites in urine by computerized GC–MS 85
Benzodiazepines
Simultaneous determination of diazepam and its active metabolites in rat serum, brain and cerebrospinal fluid by HPLC 334
Bile acids
Internal standards for quantitative GC of individual bile acids after group separation of bile acids in urine 25
Biphenyls, polychlorinated
Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex® 5000 103
Bopindolol
Column LC determination of hydrolysed bopindolol, in the picogram per millilitre range in plasma, using cartridge extraction and dual electrochemical detection 205
Butylated hydroxytoluene
Determination of butylated hydroxytoluene in plasma by HPLC 288
Caffeine
Pharmacokinetics of intravenous caffeine: comparison of HPLC and GC methods 274
Cefixime
Determination of cefixime in biological samples by reversed-phase HPLC 145
Ceftetrame
Quantitation of ceftetrame in human plasma and urine by HPLC 135
Chlorambucil
HPLC of chlorambucil analogues 315
Cimetidine
Highly sensitive determination of cimetidine and its metabolites in serum and urine by HPLC 301
Cimetidine sulphoxide
Highly sensitive determination of cimetidine and its metabolites in serum and urine by HPLC 301
Ciprofloxacin
HPLC of ciprofloxacin and its metabolites in serum, urine and sputum 153
Citric acid
Excretion of urinary acids during inverted sleep–waking rhythm 1
Codeine
Identification of opiates in urine by capillary column GC of two different derivatives 117
Cyclophosphamide
Determination of cyclophosphamide in whole blood and plasma by reversed-phase HPLC 125
Dapsone
Dapsone and monoacetyldapsone determined in serum and saliva by a sensitive HPLC method with a single extraction step 322
Desethyleneciprofloxacin
HPLC of ciprofloxacin and its metabolites in serum, urine and sputum 153
Desmethyldiazepam
Simultaneous determination of diazepam and its active metabolites in rat serum, brain and cerebrospinal fluid by HPLC 334
Desmosine
HPLC determination of desmosine and isodesmosine in tissues and its application to studies of alteration of elastin induced by atherosclerosis 53
Dextromethorphan
Direct determination of dextromethorphan and its three metabolites in urine by HPLC using a precolumn switching system for sample clean-up 340

Dextrorphan
Direct determination of dextromethorphan and its three metabolites in urine by HPLC using a precolumn switching system for sample clean-up 340
Diaminodiphenylsulphone
Dapsone and monoacetyldapsone determined in serum and saliva by a sensitive HPLC method with a single extraction step 322
Diazepam
Simultaneous determination of diazepam and its active metabolites in rat serum, brain and cerebrospinal fluid by HPLC 334
Dibenzo-*p*-dioxins, polychlorinated
Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex® 5000 103
Dibenzofurans, polychlorinated
Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex® 5000 103
3,5-Di-*tert.*-butyl-4-hydroxytoluene
Determination of butylated hydroxytoluene in plasma by HPLC 288
Dicarboxylic porphyrins
Normal-phase HPLC of free acid dicarboxylic porphyrins and hematoporphyrin derivative on silica 73
β,β-Difluorochlorambucil
HPLC of chlorambucil analogues 315
7β,12β-Dihydroxy-5β-cholan-24-oic acid
Internal standards for quantitative GC of individual bile acids after group separation of bile acids in urine 25
[3,4-Di-(4-hydroxyphenyl)-5-isoxazolyl]-acetic acid
HPLC determination of [3,4-di-(4-methoxyphenyl)-5-isoxazolyl]acetic acid and its metabolites in human plasma and urine 227
[3,4-Di-(4-methoxyphenyl)-5-isoxazolyl]-acetic acid
HPLC determination of [3,4-di-(4-methoxyphenyl)-5-isoxazolyl]acetic acid and its metabolites in human plasma and urine 227>
β,β-Dimethylchlorambucil
HPLC of chlorambucil analogues 315
N,N-Dimethyltryptamine
GC–mass spectroscopic characterisation of the psychotomimetic indolealkylamines and their in vivo metabolites 13
Disodium glyco-7α,12α-dihydroxy-5β-cholan-24-oate
Internal standards for quantitative GC of individual bile acids after group separation of bile acids in urine 25
Dithiothreitol
Simplified monitoring of reduced and oxidized dithiothreitol using HPLC 239
Elastin
HPLC determination of desmosine and isodesmosine in tissues and its application to studies of alteration of elastin induced by atherosclerosis 53
Enzymes
Assay for trimethyllysine hydroxylase by HPLC 253
Ethylmorphine
Identification of opiates in urine by capillary column GC of two different derivatives 117
Fenoprofen
HPLC quantitation of triacylglycerols containing fenoprofen from biological samples 333
1-(β-D-Glucopyranosyl)amobarbital
HPLC determination of the diastereomers of 1-(β-D-glucopyranosyl)amobarbital in urine 309
Hematoporphyrin
Normal-phase HPLC of free acid dicarboxylic porphyrins and hematoporphyrin derivative on silica 73
Heroin
Concentrations of free and conjugated morphine in blood in twenty cases of heroin-related deaths 267
Homocysteine
Determination of free and total homocysteine in human plasma by HPLC with fluorescence detection 43
Hydroxyethylvinyldeuteroporphyrin
Normal-phase HPLC of free acid dicarboxylic porphyrins and hematoporphyrin derivative on silica 73
7-Hydroxymethotrexate
Concurrent analysis of methotrexate, trimethoprim, sulphamethoxazole and their major metabolites in plasma by HPLC 328
Hydroxymethylcimetidine
Highly sensitive determination of cimetidine and its metabolites in serum and urine by HPLC 301

1-Hydroxymethyltriazolam
HPLC determination of triazolam and its metabolites in human urine 197
[3-(4-Hydroxyphenyl)-4-(4-methoxyphenyl)-5-isoxazolyl] acetic acid
HPLC determination of [3,4-di-(4-methoxyphenyl)-5-isoxazolyl] acetic acid and its metabolites in human plasma and urine 227
4-Hydroxypropranolol
Improved HPLC method for the simultaneous determination of propranolol and 4-hydroxypropranolol in plasma with fluorescence detection 294
5-Hydroxypyrazinamide
Microdetermination of pyrazinamide and its metabolites (2-pyrazinoic acid, 5-hydroxypyrazinoic acid, 5-hydroxypyrazinamide and pyrazinuric acid) in plasma and urine by LC 217
5-Hydroxypyrazinoic acid
Microdetermination of pyrazinamide and its metabolites (2-pyrazinoic acid, 5-hydroxypyrazinoic acid, 5-hydroxypyrazinamide and pyrazinuric acid) in plasma and urine by LC 217
4-Hydroxytriazolam
HPLC determination of triazolam and its metabolites in human urine 197
Indolealkylamines
GC–mass spectroscopic characterisation of the psychotomimetic indolealkylamines and their in vivo metabolites 13
Internal standards
Internal standards for quantitative GC of individual bile acids after group separation of bile acids in urine 25
Isodesmosine
HPLC determination of desmosine and isodesmosine in tissues and its application to studies of alteration of elastin induced by atherosclerosis 53
Lipids
HPLC quantitation of triacylglycerols containing fenoprofen from biological samples 33
Methotrexate
Concurrent analysis of methotrexate, trimethoprim, sulphamethoxazole and their major metabolites in plasma by HPLC 328
5-Methoxy-N,N-dimethyltryptamine
GC–mass spectroscopic characterisation of the psychotomimetic indolealkylamines and their in vivo metabolites 13
3-Methoxymorphinan
Direct determination of dextromethorphan and its three metabolites in urine by HPLC using a precolumn switching system for sample clean-up 340
[3-(4-Methoxyphenyl)-4-(4-hydroxyphenyl)-5-isoxazolyl] acetic acid
HPLC determination of [3,4-di-(4-methoxyphenyl)-5-isoxazolyl] acetic acid and its metabolites in human plasma and urine 227
Monoacetyldapsone
Dapsone and monoacetyldapsone determined in serum and saliva by a sensitive HPLC method with a single extraction step 322
6-Monoacetylmorphine
Identification of opiates in urine by capillary column GC of two different derivatives 117
β-Monomethylchlorambucil
HPLC of chlorambucil analogues 315
β-Mono(trifluoromethyl)chlorambucil
HPLC of chlorambucil analogues 315
Morphinan-3-ol
Direct determination of dextromethorphan and its three metabolites in urine by HPLC using a precolumn switching system for sample clean-up 340
Morphine
Concentrations of free and conjugated morphine in blood in twenty cases of heroin-related deaths 267
Morphine
Identification of opiates in urine by capillary column GC of two different derivatives 117
Neuropeptides
Reversed-phase HPLC–radioreceptor assay of human cerebrospinal fluid neuropeptides. Maximizing recovery of picomoles of peptides and minimizing memory 61
Nicotinamide
HPLC determination of nicotinamide in rat tissue samples and blood after extraction with diethyl ether 257
Opiates
Identification of opiates in urine by capillary column GC of two different derivatives 117
Organic acids
Excretion of urinary acids during inverted sleep–waking rhythm 1
Organochlorine pesticides

Analysis of organochlorine pesticides, poiychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex® 5000 103

Oxazepam
Simultaneous determination of diazepam and its active metabolites in rat serum, brain and cerebrospinal fluid by HPLC 334

3-Oxociprofloxacin
HPLC of ciprofloxacin and its metabolites in serum, urine and sputum 153

Pentylurofuran acid
Excretion of urinary acids during inverted sleep–waking rhythm 1

Peptides
Reversed-phase HPLC–radioreceptor assay of human cerebrospinal fluid neuropeptides. Maximizing recovery of picomoles of peptides and minimizing memory 61

Pesticides
Analysis of organochlorine pesticides, polychlorinated biphenyls, dibenzo-*p*-dioxins and dibenzofurans in human milk by extraction with the lipophilic gel Lipidex® 5000 103

Phenylacetic mustard
HPLC of chlorambucil analogues 315

Porphyrins
Normal-phase HPLC of free acid dicarboxylic porphyrins and hematoporphyrin derivative on silica 73

Propranolol
Improved HPLC method for the simultaneous determination of propranolol and 4-hydroxypropranolol in plasma with fluorescence detection 294

Propylurofuran acid
Excretion of urinary acids during inverted sleep–waking rhythm 1

Protoporphyrin
Normal-phase HPLC of free acid dicarboxylic porphyrins and hematoporphyrin derivative on silica 73

Pyrazinamide
Microdetermination of pyrazinamide and its metabolites (2-pyrazinoic acid, 5-hydroxypyrazinoic acid, 5-hydroxypyrazinamide and pyrazinuric acid) in plasma and urine by LC 217

2-Pyrazinoic acid
Microdetermination of pyrazinamide and its metabolites (2-pyrazinoic acid, 5-hydroxypyrazinoic acid, 5-hydroxypyrazinamide and pyrazinuric acid) in plasma and urine by LC 217

Pyrazinuric acid
Microdetermination of pyrazinamide and its metabolites (2-pyrazinoic acid, 5-hydroxypyrazinoic acid, 5-hydroxypyrazinamide and pyrazinuric acid) in plasma and urine by LC 217

Pyridostigmine
HPLC determination of pyridostigmine in plasma 353

Ranitidine
Automated solid-phase extraction and HPLC determination of ranitidine from urine, plasma and peritoneal dialysate 165

Salbutamol
Analysis of salbutamol enantiomers in human urine by chiral HPLC and preliminary studies related to the stereoselective disposition kinetics in man 187

Sodium glyco-7α,12α-dihydroxy-5β-cholan-24-oate
Internal standards for quantitative GC of individual bile acids after group separation of bile acids in urine 25

Sodium tauro-7β,12β-dihydroxy-5β-cholan-24-oate
Internal standards for quantitative GC of individual bile acids after group separation of bile acids in urine 25

Sulphamethoxazole
Concurrent analysis of methotrexate, trimethoprim, sulphamethoxazole and their major metabolites in plasma by HPLC 328

4-Sulphociprofloxacin
HPLC of ciprofloxacin and its metabolites in serum, urine and sputum 153

Temazepam
Simultaneous determination of diazepam and its active metabolites in rat serum, brain and cerebrospinal fluid by HPLC 334

$\beta,\beta,\gamma,\gamma$-Tetrafluorochlorambucil
HPLC of chlorambucil analogues 315

Theophylline
Improved method for the determination of theophylline in plasma and urine using HPLC 281

Thromboxane B_2
Separation of thromboxane B_2 by HPLC 246

Triacyglycerol
HPLC quantitation of triacylglycerols containing fenoprofen from biological samples 33

Triazolam
HPLC determination of triazolam and its metabolites in human urine 197
Trimethoprim
Concurrent analysis of methotrexate, trimethoprim, sulphamethoxazole and their major metabolites in plasma by HPLC 328
Trimethyllysine hydroxylase
Assay for trimethyllysine hydroxylase by HPLC 253
Uremic toxins
Ion-pair HPLC profiling of a uremic toxin fraction 263
Urinary acids
Excretion of urinary acids during inverted sleep–waking rhythm 1